

# Pleiotropy for quantitative traits: pleio

Dan Schaid, Jason Sinnwell

09 August, 2016

## Example Data

The pleio package contains a pre-made simulated dataset with multiple quantitative traits simulated from a multivariate normal distribution with common correlation structure, correlation of 0.5, and genotypes simulated based on minor allele frequency of 0.2, and assumes that the regression coefficients (beta's) for traits 2 and 3 have non-zero values, while all other traits are not associated with dose of minor allele.

Here, we load the simulated dataset and show

```
## load package and dataset
require(pleio)

## Loading required package: pleio

## Loading required package: Matrix

data(pleio.qsim)

## preview simulated data
head(y)

##          t1         t2         t3         t4         t5         t6
## [1,] -0.6795023  1.07928655 -0.3447868  0.7757427  0.2441987  0.2849724
## [2,] -0.5310806  0.06419289 -0.8963973  0.2715258 -0.4579625 -1.2279082
## [3,]  1.0244553  1.18181268  0.1977245  0.3652163  0.3497826  1.0145627
## [4,] -0.3054566 -0.49250649  1.0348276 -0.1683192  1.8157372  0.4571641
## [5,] -0.4246874 -1.57093941 -0.3505441  0.7817763 -2.4317261 -1.0474812
## [6,]  2.0213843  1.45198074  0.2430903  2.2323818  1.2128110  2.0509974

table(geno)

##  geno
##    0   1   2
##  312 170 18
```

## Sequential Pleiotropy Tests

The *pleio.q.sequential* function is a high-level way to perform sequential tests of the number of traits (and which traits) are associated with a genotype. The algorithm starts with testing the usual multivariate null hypothesis that all beta's are zero. If this test rejects, because the p-value is less than a user-specified threshold, then allow one beta to be non-zero in order to test whether the remaining beta's = 0. If the test of one non-zero beta rejects, then allow two non-zero beta's, and continue this sequential testing until the p-value for a test is greater than the specified threshold. When the sequential testing stops, one can conclude

that the final model contains the non-zero beta's, while all other beta's are inferred to be zero. Below we run two functions, *pleio.q.fit*, which performs pre-calculations on the models to be tested, and *pleio.q.sequential*, which performs the sequential pleiotropy tests.

The final result lists the indices of the non-zero beta's (the indices of the traits associated with a genotype), and the p-value that tests the fit of the final model. A p-value greater than the threshold is expected for the final model, showing that the final model fits the data well. For this example, the sequential approach stopped at 2 traits because the p-value is greater than the *pval.threshold* argument given of 0.05.

```
fit <- pleio.q.fit(y, geno)

test.seq <- pleio.q.sequential(fit, pval.threshold=.05)
test.seq

## $pval
## [1] 0.2744734
##
## $index.beta
## [1] 2 3
```

## Equivalent Steps to Sequential Fit

The sequential steps above can be performed with more user control using *pleio.q.test*, with *count.nonzero.beta* as the number of non-zero betas for the null hypothesis. The result of *pleio.q.test* contains the global test statistic, degrees of freedom (df), p-value for testing the model, the index of the non-zero beta's in the model, and a data.frame called “tests” that contains the tests performed for the null hypothesis models (i.e., the indices of the non-zero beta's and the corresponding statistic, tk, for each model). For *m* traits, and *k* = *count.nonzero.beta*, there are m-choose-k models in the set that are considered in the null hypothesis, and the minimum tk test statistic over the set provides the global test statistic reported.

```
test0 <- pleio.q.test(fit, count.nonzero.beta = 0)
test0

## $stat
## [1] 37.08576
##
## $pval
## [1] 1.694389e-06
##
## $df
## [1] 6
##
## $index.nonzero.beta
## [1] 0
##
## $tests
##   index.1      tk
## 1       0 37.08576

test1 <- pleio.q.test(fit, count.nonzero.beta = 1)
test1
```

```

## $stat
## [1] 23.53879
##
## $pval
## [1] 0.000266202
##
## $df
## [1] 5
##
## $index.nonzero.beta
## [1] 3
##
## $tests
##   index.1      tk
## 1      1 32.44700
## 2      2 24.26597
## 3      3 23.53879
## 4      4 28.02627
## 5      5 37.01172
## 6      6 36.32446

test2 <- pleio.q.test(fit, count.nonzero.beta = 2)
test2

```

```

## $stat
## [1] 5.1274
##
## $pval
## [1] 0.2744734
##
## $df
## [1] 4
##
## $index.nonzero.beta
## [1] 2 3
##
## $tests
##   index.1 index.2      tk
## 1      1      2 21.43943
## 2      1      3 21.50900
## 3      1      4 20.53175
## 4      1      5 32.01780
## 5      1      6 31.14303
## 6      2      3 5.12740
## 7      2      4 17.74535
## 8      2      5 24.14999
## 9      2      6 24.26479
## 10     3      4 17.37530
## 11     3      5 23.31140
## 12     3      6 23.24654
## 13     4      5 27.59217
## 14     4      6 25.76981
## 15     5      6 36.13547

```